

PRESCRIPTION TREATMENT[®] brand

ULTRACIDE[®]

Pressurized Flea IGR & Adulticide

Contains Nylar* Insect Growth Regulator

Formula 1

KILLS: Active flea infestations and prevents infestations from developing. Prevents reinfestation for 7 months.

FOR USE IN: Apartments, Automobiles, Homes, Hospitals, Hotels, Kennels, Motels, Offices, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks), Warehouses, Utilities, Veterinary Clinics and Other Commercial and Industrial Buildings.

ACTIVE INGREDIENTS:

Pyriproxyfen	0.100%
Pyrethrins	0.050%
N-octyl bicycloheptene dicarboximide**	0.400%
Permethrin†	0.400%
Related Compounds	0.035%

OTHER INGREDIENTS: 99.015%

* Nylar and MGK - Reg. TM of McLaughlin Gormley King, Inc.

** MGK 264, Insecticide Synergist

† Cis-trans isomers ratio: Max. 55% (+ or -) cis and Min. 45% (+ or -) trans

EPA Reg. No.: 499-404 EPA Est. No.: 499-MO-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional precautionary statements.

GENERAL INFORMATION

SHAKE WELL BEFORE USE

Ultracide[®] kills fleas and prevents flea infestations for a full season. Adult fleas may be seen in the treated areas when brought in on infested animals or when adults emerge from pupal cases, however a population will not develop in the treated area.

In transportation equipment, certain plastics may be whitened or dulled. Apply only to carpets and cloth upholstered areas, or test an inconspicuous area before treating.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Use Ultracide[®] at the rate of 20 oz. for up to 2,625 sq. ft. (10 oz. for up to 1,300 sq. ft). Carpeted areas may be vacuumed before treatment and after treatment has dried.

Treat infested areas or areas which could be infested; these include rugs, carpets, upholstered furniture, pet beds and pet resting areas. Hold can at arms length and direct spray toward the area to be treated. Use a sweeping motion to apply product and back away from treated area while holding the can 36 inches away from the surface being treated. An area of 80 - 100 sq. ft. can be treated in approximately 10 seconds with Ultracide[®]. When treating upholstered furniture, treat under cushions and areas where flea development occurs.

Cover aquariums and fish bowls and remove birds from area prior to treating. Treated areas should be vacated during application. EXCEPT FOR APPLICATOR DO NOT PERMIT HUMANS OR PETS TO CONTACT TREATED SURFACES UNTIL THE SPRAY HAS DRIED. Do not treat pets with this product.

NOTICE: Ultracide[®] has been tested on several types of carpet, fabric and other household furnishings without adverse affects. In a few instances waxed surfaces have been whitened or dulled. Holding the can 36 inches away from the target surface normally prevents any whitening from occurring. Because of the wide variety of floor types and finishes, treat a small inconspicuous area before treating the entire area. Avoid contact with antique finishes.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.



WHITMIRE MICRO-GEN
RESEARCH LABORATORIES, INC.

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.



ULTRACIDE® Pressurized Flea IGR & Adulticide

CONTAINER DISPOSAL: This container may be recycled in the few but growing number of communities where steel aerosol can recycling is available. Before offering for recycling, empty the can by using the product according to the label (DO NOT PUNCTURE!). If recycling is not available, wrap the container and discard in the trash.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



PRECAUTIONARY STATEMENTS

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Do not breathe vapors or spray mist. Avoid contact with skin or eyes. In case of contact, flush with plenty of water. Wash with soap and warm water after use. Obtain medical attention if irritation persists. Avoid contamination of food or feedstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and birds. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift from treated areas may be hazardous to organisms in adjacent aquatic sites. Apply this product only as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS

Extremely flammable. Contents under pressure. Keep away from fire, sparks, and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

A PRESCRIPTION TREATMENT® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682

© 2000 Whitmire Micro-Gen Research Laboratories, Inc.

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.

Revised: 6/00